

## FRUIT JUICE NUTRITION & HEALTH – IFU SCIENTIFIC REVIEW

### General Dietary Recommendations

The lack of adequate consumption of fruit and vegetables has become a worldwide dietary concern since fruits and vegetables play a pivotal role in attaining and maintaining good health. Decades of research have found that fruits and vegetables are crucial dietary components that can help reduce the risk for numerous chronic diseases which, in many cases, have been shown to be initiated by long term inflammation (Holt et al. 2009). The World Health Organization (WHO) as a cosponsor of the global 5+ a day program promotes the inclusion of at least five servings a day of fruit and vegetables as an essential element in a healthy diet (WHO, 2004). Practical aspects of the program can be found at the CDC (Centers for Disease Control and Prevention) web site (<http://www.fruitsandveggiesmatter.gov>). One hundred percent (100%) fruit juices along with whole fruits and vegetables are deemed to be an integral part of the 5 + a day program. The USDA, in the description of the Food Pyramid, states that 100% fruit juice counts as part of the fruit group (<http://www.mypyramid.gov/pyramid/fruits.html>), and the American Academy of Pediatrics suggests that, although fruit juices should be consumed with moderation, 6 oz of juice can count toward a serving of fruit (Pediatrics. 2001). Another report by Nicklas et al (2008) showed that American children consume only 4.1 fl oz of juice daily, which contributed to 58 kcal or 3.3% of total energy intake.

### Fruit and Fruit Juice

According to the CODEX General Standard for Fruit Juices and Nectars “*fruit juices have the essential physical, chemical, organoleptical, and nutritional characteristics of the fruit(s) from which it comes* (CODEX STAN 247-2005). Properly extracted juices are very similar to the fruit; they contain most substances which are found in the original ripe and sound fruit from which the juice is made. A fruit juice is made from the whole fruit (edible parts) and does not contain more sugar than the corresponding fruit. Recently Ruxton et al (2006) reviewed the literature comparing health benefits of fruits and fruit juices and concluded that there were no significant differences. The similarities between the composition of fruits and fruit juices were also pointed out by Landon (2007).

Juices and juice-puree mixtures (also known as “smoothies”) are more convenient to consume, and have in general a longer shelf-life than fresh fruit. Therefore, moderate intake of juices along with fruit is deemed to be completely appropriate by the 5+ a day programme. Fruit juice is free of chemical preservatives and other additives. Only in a few cases, for example cloudy apple juice, the addition of ascorbic acid (vitamin C) is used for the prevention of browning. As a mirror image of the fruit, it contains essentially all substances which are found in the original fruit which must be ripe and healthy. It is the major task of modern food technology to transfer the valuable fruit components into the juice and to produce stable products by physical means.

A fruit juice is always a 100 % product and should not be confused with soft drinks or other refreshing drinks.

## **Fruit juice health promoting components**

Fruits and vegetables form a versatile and complex substance group category of foods. The relevant substance groups are carbohydrates, acids, minerals, polyphenols (tannins) including the colourful anthocyanins, water-soluble vitamins, amino acids, aroma compounds, carotenoids, fibers and other bioactive substances. During processing, they are essentially transferred into the pressed juice or into the puree.

Juices are low in compounds such as sodium and fat which are believed to have negative health impacts when ingested in large amounts. Conversely, juices contain a variety of beneficial micronutrients, including significant quantities of minerals, like potassium as the major mineral (Dillon, 1995), followed by calcium and magnesium. Many trace elements of fruits are also found in the corresponding fruit juices. The B-vitamin folate is present in orange, grapefruit, pineapple juices and some other tropical juices. Vitamin C, although sometimes added to fruit juices, is found in significant amounts in different fruit juices. Examples are orange juice, grapefruit juice, black currant juice, strawberry juice, or acerola juice. These properties have given rise to most of the health claims that have been approved by supervising organizations. Health claims have been formulated under Article 13 of the EC Regulation on nutrition and health claims (2009) and by the FDA in 21 Code of Federal Regulations Part 101. In recent years, the beneficial phytochemicals present in all the major fruits and fruit juices have been characterized and a database on the polyphenol food content is now available online (<http://www.phenol-explorer.eu/>, 2009).

Apples and cloudy apple juice contain quercetin, chlorogenic and other phenolic acids as well as phloridzin and phloretin xyloglucoside (Soler et al, 2009). The two last mentioned substances belong to the group of dihydrochalcones which are typical for apple products and seem to play a positive role in the regulation of blood glucose level.

Berries are rich in anthocyanins (cyanidin, pelargonidin) and flavonoids (kaempferol and quercetin derivatives) (Szajdek and Borowska, 2008, Seeram, 2008). Among those, cranberries have been particularly studied (Neto 2007).

Citrus contains vitamin C, Thiamin, folate, flavanones (hesperidin, naringin), carotenoids such as beta-carotene, alpha-carotene, beta-cryptoxanthin, lutein zeaxanthin, and lycopene in red grapefruit, (Baghurst, 2003; Benavente-García and Castillo, 2008; Marti et al, 2009), finally limonoids are found in grapefruits (Manners, 2007).

Pomegranate juice is rich in ellagitannins, like punicalagin (Basu and Penugonda, 2009). These substances are antioxidative and antiinflammatory. Other ellagitannins are found in berry juices and nectars made from blackberry, raspberry, and strawberry. Grape juice is well known for the presence of resveratrol, flavonoids and anthocyanins (Pezzuto et al, 2009; Marquez et al, 2009; Iriti and Faoro, 2009) and tomato juice<sup>1</sup> is a major source of lycopene (Lee et al, 2009).

In addition, juices may serve as carriers for added nutrients such as calcium, vitamin D and phytosterols that may not be inherent in the fruit itself.

## **Fruit Juice Phytochemical Properties**

In the last ten years, the mechanisms responsible for the health benefits of fruit juices have been considerably investigated. The biological activities cannot be solely explained

---

<sup>1</sup> *Vegetable juices are assumed to be within the term 'fruit juice' and their properties are referred to without distinguishing them as a vegetable source of juice.*

by antioxidant effects. Most of the antioxidant benefits of food based on chemical oxidation reactions have now been shown to be unfounded (EFSA Journal 2010; 8(2):1489). Antioxidant effects occur through the up or down regulation of specific enzymes. It has been shown that phytochemicals act through the modulation of signal cascades in the human body, most often starting at the genes level. Based on these interactions signals are transferred into the cells leading to activation or deactivation of metabolic pathways. (Serafini, 2009; Koltover, 2009; Crozier et al 2009).

The question of the mode of action of phytochemicals was raised as early as 2004 by Azzi et al. for vitamin E. The lack of relationship between in vitro antioxidant activity of phytochemicals and their physiological properties was examined by most groups involved in the health benefits of these compounds. (Williams et al, 2004; Cerda et al 2004; Scalbert et al, 2005; Sies, 2007; Stevenson and Hurst, 2007) Similar processes have now been reported for lycopene (Erdman et al. 2009) and even vitamin C (Wu et al, 2007; Kelly et al, 2008; Maeng et al, 2009). It is now accepted that phytochemicals, once ingested, are modified and metabolized in the intestinal tract. Parts of these phytochemicals and of their metabolites are absorbed into the blood stream, modified again in the liver and other organs. Finally, these metabolites act at the gene level and start a series of cascading reactions promoting physiological changes. Recent studies have shown that following orange juice ingestion, genes are affected much sooner than cardiovascular health markers (Morand et al. 2011a,b). This has led to the development of new analytical procedures to examine the effects of fruit juices and other foods, based on genomics, proteomics and metabolomics (Ovesna, 2008; Scalbert and Knasmüller, 2008; Fardet et al, 2008; Steiner et al, 2008; Mauray et al 2010). It should be kept in mind that, in many cases, the end reaction of these pathways is an antioxidant effect occurring through the modulations of enzymes such as catalase, NADPH oxidase, superoxide dismutase, and glutathione peroxidase. This has led some supervising authorities to question the meaning of the word "antioxidant" that they have qualified as ambiguous (Afssa – Saisine n° SA-0266, 2006). There is therefore a need to redefine the meaning of the words used to define the beneficial activities of fruit juice phytochemicals.

### **Health Benefits of Fruit Juices**

The health benefits of fruits and fruit juices have been reviewed by Landon (2007). The high potassium and low sodium characteristic of most juices help maintain a healthy blood pressure, furthermore the lack or near absence of fat in fruit juices is beneficial for the cardiovascular system. (Delichatsios and Welty, 2005) The fortification of juices with calcium (Andon, 1996) and phytosterol (Devaraj et al, 2004) provide some supplemental bone and cardiovascular benefits. Recently, several reviews have summarized the health benefits of fruit polyphenols (Spencer, 2010; Chong et al, 2010; Gonzalez-Gallego et al, 2010; Hardcastle et al, 2010).

Vitamins have a special role since they are essential for life and most are not produced by the body. Vitamin C (ascorbic acid), naturally present or added to most juices, is necessary for the body to form collagen, cartilage, muscle, and blood vessels, and aids in the absorption of iron. The enzymatic and non enzymatic functions of vitamin C were reviewed by Levine et al (1993). Its role as an antioxidant has been extensively examined, however many vitamin C effects appear to be due to its role as a coenzyme in many biochemical reactions (Levine et al, 1993). More recently, the influence of vitamin C in gene modulation and biochemical pathways modifications has been shown, particularly in blood vessel endothelium (Wu et al, 2007) and atherosclerosis (Frikke-

Schmidt and Lykkesfeldt, 2009). Among the vitamins found in fruit juices, folate from citrus and pineapple is essential for the prevention of spina bifida (Bell and Oakley, 2009) and premature birth ( Bukowsky et al, 2009). It also helps in maintaining a low level of the amino acid homocysteine, a marker of inflammation, that has been associated with a higher risk for heart disease, stroke, and heart failure ( Sánchez-Moreno et al, 2009).

The health benefits of minerals, vitamins, and micronutrients have been well characterized but many beneficial properties of juices have been shown to come from phytochemicals, mainly polyphenols, carotenoids and limonoids. It should be pointed out that data on the health benefits of fruit juices are still fragmentary and that many studies have been done on cell cultures and animal models. Efforts are to be made to perform extended clinical studies with doses of juices similar to those ingested in normal life. If health claims are to be accepted, it will be necessary to determine the bioavailability of the main fruit juice phytochemicals and to define some type of RDI values (Williamson and Holst, 2008; Holst and Williamson, 2008).

Most of the diseases that seem to be targeted by phytochemicals are those induced by chronic inflammation. This process, which had been called the silent killer, is an attempt by the organism to remove injurious stimuli and to initiate healing. The diseases that may be initiated by chronic inflammation include aging diseases such as Alzheimer's disease (Granic et al, 2009; Kanapuru, 2009), diabetes, insulin resistance (King, 2008; Blüher, 2008) and cardiovascular disease, particularly atherosclerosis (Bucova et al, 2008). In addition, bone diseases such as osteoporosis and arthritis (Hardy and Cooper, 2009), cognitive functions and brain diseases (Wärnberg et al, 2009) and some forms of cancer (Gonda et al, 2009) may be induced by chronic inflammation.

### **Cardiovascular Health**

The area of cardiovascular diseases is, so far, the one that appears to show the most promising potential for the beneficial effects of fruit juices (Duthie et al, 2006; Leifert and Abeywardena, 2008; Ross, 2009; Grassi et al, 2009; Dalgård et al, 2009; Chong et al, 2010).

The formation of a blood clot in the circulatory system (thrombosis) can lead to disturbance in the blood supply resulting in embolism and stroke. Several fruit juices seem to be able to limit blood clot formation by preventing platelets from agglutinating in the blood vessels (Freedman et al, 2001; Mattiello et al, 2009).

Atherosclerosis is the condition in which an artery wall thickens as the result of a accumulation of fatty material. This build up, called plaque, hardens and narrows arteries. The blood flow is reduced and can lead to heart attack and stroke. Fruit juices have been shown to act at the various levels of the processes leading to atherosclerosis. Several years ago, it was shown that fruit juices can increase the level of high density lipoproteins (HDL), the lipids disposed of in the liver (good lipids) and decrease the formation and oxidation of low density lipoproteins (LDL) that are deposited in the blood vessels (bad lipids) (Gorinstein et al, 2004, 2006). Although, preventing the oxidation of LDL may play an important role in the beneficial effects of fruit juices (Aviram et al, 2002), it appears that the effects of fruit juice phytochemicals is complex and involves gene modulation. Fruit juice components have been shown to act at every level of the blood lipid process from cholesterol synthesis to the formation of lipoproteins (LDL, HDL). For about ten years it has been known that naringin and hesperidin inhibit the first

enzyme in the biosynthesis of cholesterol (HMG-Co Reductase), the same enzyme targeted by the statin class of drugs (Bok et al, 1999). Nahmias et al (2008) showed that naringin inhibits the transcription of HMG-Co reductase, the activity of microsomal triglyceride transfer protein (MTP) and the transcription of acyl-coenzyme A: cholesterol acyltransferase 2 (ACAT2) the enzyme which in the final phase of LDL production attaches cholesterol to the lipoproteins. Similarly, naringin (Mulvihill et al, 2009) and anthocyanins from berries (Qin et al, 2009) have a beneficial effect on lipoprotein profiles by decreasing LDL-cholesterol and increasing HDL-cholesterol concentrations. Apple polyphenols (Lam et al, 2008) may act by inhibiting cholesterol ester transfer protein (CETP). Morin et al (2008) showed that a reduction of plasma cholesterol by citrus flavonoids is associated with a modulation of the expression of the LDL receptor (LDLR) gene.

Another signaling molecule affected by fruit juices is nitric oxide (NO). The endothelium (inner lining) of blood vessels uses nitric oxide to signal the surrounding smooth muscle to relax, thus resulting in vasodilation and increasing blood flow. The proper level of NO is largely modulated by enzymes such as endothelial nitric oxide synthase (eNOS), and nitric oxide oxidase. Fruit juices play a major role in the maintenance of NO levels. This has been shown to occur with many juices (George et al, 2009) including grape (Ekshyyan, et al 2007), pomegranate (de Negrís et al 2005) and citrus juices (Morand et al, 2011a,b). These beneficial changes have been noted at various levels of the cardiovascular system including blood pressure (Reshef et al, 2005; Morand et al, 2011a,b).

### **Bone Health**

Bone health is largely the result of an equilibrium between osteoclast cells destroying bone and osteoblasts building it. Several fruit juice phytochemicals, mainly polyphenols and carotenoids, have been shown to have a positive influence on bone health and particularly the bone mineral density of post-menopausal women (Trzeciakiewicz et al, 2009, 2010). Positive activity has been reported for citrus (Deyhim et al, 2008; Mandadi et al, 2009) and pomegranate (Mori-Okamoto et al, 2004). Citrus juice hesperidin and naringin may act through bone morphogenetic proteins (BMPs) pathways that induce the formation of bone and cartilage (Wong and Rabie, 2006; Trzeciakiewicz et al, 2009; Habauzit, et al 2007, 2009; Chiba et al 2007; Horcajada et al, 2008). Fruit juice carotenoids,  $\beta$ -cryptoxanthin,  $\beta$ -carotene and lycopene may also improve bone health by preventing bone destruction by osteoclasts (Sugiura et al, 2008; Sahni et al, 2009 a,b). It should also be mentioned that citrus and pomegranate may have a positive effect on arthritis (Murakami et al, 2007; Shukla et al 2008; Hadipour-Jahromy and Mozaffari-Kermani, 2010).

### **Brain health, cognition and aging**

Many reports have shown that fruit juices may play an important role in maintaining cognition, limiting brain aging and possibly slowing the progress of Alzheimer's disease (Dai et al, 2006). The beneficial effects of grape juice (Joseph et al, 2009; Wang et al 2008), berries (Shukitt-Hale et al, 2008, 2009; Willis et al 2008) and citrus (Datla et al 2001) have been examined. The ability of juice compounds, particularly flavonoids, to cross the barrier protecting the brain (blood brain barrier) is at the origin of the beneficial activity of these compounds (Youdim, 2003). Spencer's group reviewed the

neuroprotective properties of dietary flavonoids (Spencer 2008, 2009a, 2010). Firstly, they promote cerebral vascular blood flow. Secondly, they interact with neuronal signaling cascades leading to an inhibition of cell death and to a promotion of neuronal differentiation. As a result, they prevent deterioration or even improve cognitive performance (Macready et al 2009; Spencer, 2009b; Vafeiadou et al 2009; Harrison and May, 2009). Vitamin C can also reach the brain and ascorbate is proposed as a neuromodulator of neurotransmitters, thus vitamin C may have potential therapeutic roles against ischemic stroke, Alzheimer's disease, Parkinson's disease, and Huntington's disease (Harrison and May, 2009).

### **Cancer and inflammation**

Several epidemiological studies have shown that fruit juices may have a beneficial role in preventing the development of cancer (Cutler et al, 2008; Wu, et al, 2009, Kyle et al, 2009). Many juice phytochemicals, polyphenols, carotenoids and limonoids may influence mechanisms relevant for cancer prevention. These include antimutagenic activity, control of angiogenesis, anti-inflammatory mechanisms, modulation of signal transduction pathways. Positive results have been associated with most juices, including apple juice (Gerhaeuser 2008, Veeriah et al, 2008) grapes and grape juice (Iriti and Faoro 2009). Anthocyanins from various berry juices (Hochman et al, 2008; Thomasset et al, 2009; Matsunaga, 2010), citrus flavonoids (Benavente-García and Castillo 2008) and limonoids (Poulose et al, 2006) may also have potential anticarcinogenic activities. Among those, the control of chronic inflammation may be very important (González-Gallego et al, 2010).

In a different area related to inflammation, cranberry juice has long been associated with a reduction in urinary infections. Anthocyanidin and proanthocyanidin may inhibit the adhesion of uropathogens (e.g. uropathogenic *E. coli*) to the uroepithelium, thus impairing colonization and subsequent infection (Guay, 2009).

### **Skin health**

In recent years, studies have shown that juices and their phytochemicals can improve skin health (Cosgrove et al, 2007) and have sometimes been referred to as providing "Beauty from within". Vitamin C has long been known to maintain skin collagen (Sheretz and Goldsmith 1991). A beneficial effect of vitamin C on skin, starting at the gene expression level has been revealed (Arai et al 2009, Duarte et al 2009). Flavonoids have been shown to improve skin microcirculation (Neukam et al, 2007) and collagen formation (Stipcevic et al, 2006; Bae et al, 2009). Carotenoids have also been shown to improve skin health (Stahl and Sies, 2007).

### **Body weight and insulin resistance**

Excessive consumption of any calorie source is likely to lead to obesity. For example, Faith et al (2006) showed that the consumption of very high levels of fruit juice could have a negative effect (24-30 fl oz, 709-889 ml). Concern has been expressed that fruit juice sugar can contribute to weight gain, especially for children. However, O'Connor et al (2006) reported that on average, preschool children drank less than 177 ml/day 100% fruit juice. Similarly, Nicklas et al (2008) reported that the mean daily juice consumption was 4.1 fl oz, which contributed a mean intake of 58 kcal (3.3% of total energy intake).

On average, children consumed less than the maximum amounts of 100% juice recommended by the American Academy of Pediatrics. The Academy recommends limiting fruit juice consumption to 4 – 6 ounces/day for children 1 to 6 years old and 8 – 12 ounces or 2 servings/day for children 7 – 18 years old (Pediatrics, 2001).

Several studies have looked at the effects of drinks by grouping sodas and 100% fruit juice, and examining populations of various sizes, therefore providing skewed and confusing data (Palmer et al, 2008). These questions were reviewed by Nicklas et al (2008). O'Neil and Nicklas (2008) evaluated the relationship between consumption of 100% fruit juice and bodyweight among children and adolescents and found that there is no systematic association between consumption of 100% fruit juice and overweight in children or adolescents. The findings of this analysis support previous studies by Skinner et al (1999, 2001) and Newby et al (2004) that showed no significant correlation between weight change and the consumption of 100% fruit juice, fruit drinks, milk, soda, or diet soda in preschool-aged children. It has been suggested that the amount of sugar being absorbed can be monitored by using the Glycemic Index (GI). This concept has led to a reassessment of the role of sugars in juices within the diet. Slowly absorbed foods have a low GI rating while foods that are more quickly absorbed will have a higher rating (Foster-Powell et al, 2001; Mendosa, 2003). However, the glycemic index tells you how quickly a carbohydrate-containing food turns into sugar, but it doesn't tell you how much carbohydrate is in a serving of a food (Vrolix and Mensink, 2010). In many cases, the use of the glycemic load which considers both the glycemic index and the amount of carbohydrate in a food would be more appropriate. The carbohydrate in carrots, for example, has a high GI, but carrots are low in carbohydrate compared to other foods, so carrots glycemic load is relatively low (Venn and Green, 2007).

Study of fruit juice intake based only on calorie calculations ignores several important points. It has been shown that fruit juices contain compounds that limit or prevent insulin resistance by fruit juices. Yoshida et al (2007) showed that fruit juice consumption was inversely associated with fasting plasma glucose. A reduction of insulin resistance, oxidative stress and inflammation were reported after ingestion of several fruit juices including grapefruit juice (Yao et al, 2004), grapefruit naringin (Kannappan and Anuradha, 2009), orange juice (Ghanim et al 2007), cranberry juice (Wilson et al, 2008), and blueberry, (DeFuria et al, 2009). These results were recently confirmed by Wu et al (2009) who showed that flavonoids can attenuate the expression of glucose induced inflammatory cytokines. Alcada et al (2009) also showed that orange juice may increase the production of somatostatin, an inhibitor of insulin secretion. The authors concluded that fruit juices are a good source of the sugars needed by the body, and also provide phytochemicals with a wide range of health benefits.

## **Dental health**

Several reports have postulated that fruit juices can affect dental health, promote caries and dissolve enamel. Reports of these effects appeared as early as 1954 (Thomas). These studies have been performed either in vitro or under, very long, non natural conditions. (Willershausen et al, 2008; Ren et al 2009; Ehlen et al, 2008). Lim et al (2008) reported that children with a high consumption of soft drinks are at a higher risk of developing dental caries relative to those consuming milk and 100 percent fruit juice, and that damage can be avoided by regular use of a fluoridated dentifrice. No measurable association of intake of 100% fruit juice, milk, tea and some other drinks with the prevalence of tooth loss were reported by Tanaka et al (2008).

Hooper et al (2007) suggested that the sodium hexametaphosphate containing paste could be used to provide significant erosion protection in susceptible individuals. While possible detrimental effect of juice sugar and acidity can easily be eliminated by proper hygiene, recent studies have revealed that juice polyphenols have a beneficial effect on dental health (Hannig et al 2009; Ferrazzano et al, 2009). Furthermore, calcium added to juice can also add another beneficial effect (Davis et al, 2007).

## **Conclusion**

Fruit juices contain nutrients, minerals, trace elements, vitamins and phytochemicals that have been shown to have many health benefits. Although studies are still fragmentary and need to be expanded, particularly in the clinical area, juices appear to be most active on diseases related to chronic inflammation, cancer, heart and bone diseases, problems related to cognition and aging, and possibly insulin resistance. The mode of action of those fruit juice compounds in most cases seems to be by modulating gene activity. Fruit juices, consumed in moderation as part of a balanced diet, offer both: health and disease risk reduction properties. Furthermore, to identify fruit juices as inadvisable in the context of obesity and dental health, would deny the consumer a perfectly healthy and nutritious food, and be completely contrary to the evidence noted in the scientific community.

Fruit juices have a positive effect on the human health, but like any plant foods they are not drugs and cannot and should not act as such.

## FRUIT JUICE NUTRITION & HEALTH – IFU SCIENTIFIC REVIEW

### REFERENCES

- Agence française de sécurité sanitaire des aliments- Afssa – Saisine n° (2006)-SA-0266.
- Alçada MN, Monteiro R, Calhau C, Azevedo I. Orange juice vs soft drinks. *Arch Intern Med.* (2009), 169:315.
- American Academy of Pediatrics: The use and misuse of fruit juice in pediatrics. Committee on Nutrition. *Pediatrics.* (2001), 107:1210-1213.
- Andon MB, Peacock M, Kanerva RL, De Castro JA. Calcium absorption from apple and orange juice fortified with calcium citrate malate (CCM). *J Am Coll Nutr.* (1996), 15:313-316.
- Arai KY, Sato Y, Kondo Y, Kudo C, Tsuchiya H, Nomura Y, Ishigami A, Nishiyama T. Effects of vitamin C deficiency on the skin of the senescence marker protein-30 (SMP30) knockout mouse. *Biochem Biophys Res Commun.* (2009) 385:478-483
- Aviram M, Dornfeld L, Kaplan M, Coleman R, Gaitini D, Nitecki S, Hofman A, Rosenblat M, Volkova N, Presser D, Attias J, Hayek T, Fuhrman B. Pomegranate juice flavonoids inhibit low-density lipoprotein oxidation and cardiovascular diseases: studies in atherosclerotic mice and in humans. *Drugs Exp Clin Res.* (2002), 28:49-62.
- Azzi A, Davies KJ, Kelly F. Free radical biology - terminology and critical thinking. *FEBS Lett.* (2004), 558:3-6.
- Bae JY, Lim SS, Kim SJ, Choi JS, Park J, Ju SM, Han SJ, Kang IJ, Kang YH. Bog blueberry anthocyanins alleviate photoaging in ultraviolet-B irradiation-induced human dermal fibroblasts. *Mol Nutr Food Res.* (2009), 53:726-38.
- Baghurst K. The Health Benefits of Citrus Fruits. *Health Sciences & Nutrition.* CSIRO. (2003).
- Basu A, Penugonda K. Pomegranate juice: a heart-healthy fruit juice. *Nutr Rev.* (2009), 67:49-56.
- Bell KN, Oakley GP Jr. Update on prevention of folic acid-preventable spina bifida and anencephaly. *Birth Defects Res A Clin Mol Teratol.* (2009),85:102-107.
- Benavente-García O, Castillo J. Update on uses and properties of citrus flavonoids: new findings in anticancer, cardiovascular, and anti-inflammatory activity. *J Agric Food Chem.* (2008), 56:6185-6205.
- Blüher M. The inflammatory process of adipose tissue. *Pediatr Endocrinol Rev.* (2008), 6:24-31.
- Bok SH, Lee SH, Park YB, Bae KH, Son KH, Jeong TS, Choi MS. Plasma and hepatic cholesterol and hepatic activities of 3-hydroxy-3-methyl-glutaryl-CoA reductase and acyl CoA: cholesterol transferase are lower in rats fed citrus peel extract or a mixture of citrus bioflavonoids. *J Nutr.* (1999), 129:1182-1185.
- Bucova M, Bernadic M, Buckingham T. C-reactive protein, cytokines and inflammation in cardiovascular diseases. *Bratisl Lek Listy.* (2008), 109:333-340.
- Bukowski R, Malone FD, Porter FT, Nyberg DA, Comstock CH, Hankins GD, Eddleman K, Gross SJ, Dugoff L, Craigo SD, Timor-Tritsch IE, Carr SR, Wolfe HM,

- D'Alton ME. Preconceptional folate supplementation and the risk of spontaneous preterm birth: a cohort study. *PLoS Med.* (2009).
- Cerdá B, Espín JC, Parra S, Martínez P, Tomás-Barberán FA. The potent in vitro antioxidant ellagitannins from pomegranate juice are metabolised into bioavailable but poor antioxidant hydroxy-6H-dibenzopyran-6-one derivatives by the colonic microflora of healthy humans. *Eur J Nutr.* (2004), 43:205-220.
- Chiba H, Uehara H, Ishimi Y, Suzuki K, Kim Y, Matsumoto A. Hesperidin, a citrus flavonoid, prevents bone loss in ovariectomized aged mice. *Third International Conference on Polyphenols and Health* (2007).
- Chong MF, Macdonald R, Lovegrove JA. Fruit polyphenols and CVD risk: a review of human intervention studies. *Br J Nutr.* (2010)104 Suppl 3:S28-39.
- Codex general standard for fruit juices and nectars (CODEX STAN 247-2005)
- Cosgrove MC, Franco OH, Granger SP, Murray PG, Mayes AE. Dietary nutrient intakes and skin-aging appearance among middle-aged American women. *Am J Clin Nutr.* (2007), 86:1225-31.
- Crozier A, Jaganath IB, Clifford MN. Dietary phenolics: chemistry, bioavailability and effects on health. *Nat Prod Rep.* (2009), 26:1001-1043.
- Cutler GJ, Nettleton JA, Ross JA, Harnack LJ, Jacobs DR Jr, Scrafford CG, Barraja LM, Mink PJ, Robien K. Dietary flavonoid intake and risk of cancer in postmenopausal women: the Iowa Women's Health Study. *Int J Cancer.* (2008), 123:664-671.
- Dai Q, Borenstein AR, Wu Y, Jackson JC, Larson EB. Fruit and vegetable juices and Alzheimer's: the Kame Project. *Am J Med.* (2006), 119:751-9.
- Dalgård C, Nielsen F, Morrow JD, Enghusen-Poulsen H, Jonung T, Hørder M, de Maat MP. Supplementation with orange and blackcurrant juice, but not vitamin E, improves inflammatory markers in patients with peripheral arterial disease. *Br J Nutr.* (2009),101:263-9.
- Datla KP, Christidou M, Widmer WW, Rooprai HK, Dexter DT. Tissue distribution and neuroprotective effects of citrus flavonoid tangeretin in a rat model of Parkinson's disease. *Neuroreport.* (2001), 12:3871-5.
- Davis RE, Marshall TA, Qian F, Warren JJ, Wefel JS. In vitro protection against dental erosion afforded by commercially available, calcium-fortified 100 percent juices. *J Am Dent Assoc.* (2007), 138:1593-1598.
- De Nigris F, Williams-Ignarro S, Lerman LO, Crimi E, Botti C, Mansueto G, D'Armiento FP, De Rosa G, Sica V, Ignarro LJ, Napoli C. Beneficial effects of pomegranate juice on oxidation-sensitive genes and endothelial nitric oxide synthase activity at sites of perturbed shear stress. *Proc Natl Acad Sci U S A.* (2005),102:4896-4901.
- DeFuria J, Bennett G, Strissel KJ, Perfield JW 2nd, Milbury PE, Greenberg AS, Obin MS. Dietary blueberry attenuates whole-body insulin resistance in high fat-fed mice by reducing adipocyte death and its inflammatory sequelae. *J Nutr.* (2009), 139:1510-1516.
- Delichatsios HK, Welty FK. Influence of the DASH diet and other low-fat, high-carbohydrate diets on blood pressure. *Curr Atheroscler Rep.* (2005), 7:446-454.

- Devaraj S, Jialal I, Vega-López S. Plant sterol-fortified orange juice effectively lowers cholesterol levels in mildly hypercholesterolemic healthy individuals. *Arterioscler Thromb Vasc Biol.* (2004), 24:e25-8.
- Deyhim F, Mandadi K, Faraji B, Patil BS. Grapefruit juice modulates bone quality in rats. *J Med Food.* (2008),1:99-104.
- Dillon A, Fruit juice profiles. In “Methods to detect adulteration of fruit juice beverages”. Nagy S, Wade RL. eds. (1995), AGscienceAuburndale, Fl.. Appendix II 359-433.
- Duarte TL, Cooke MS, Jones GD. Gene expression profiling reveals new protective roles for vitamin C in human skin cells. *Free Radic Biol Med.* (2009),46:78-87.
- Duthie SJ, Jenkinson AM, Crozier A, Mullen W, Pirie L, Kyle J, Yap LS, Christen P, Duthie GG. The effects of cranberry juice consumption on antioxidant status and biomarkers relating to heart disease and cancer in healthy human volunteers. *Eur J Nutr.* (2006), 45:113-122.
- Ehlen LA, Marshall TA, Qian F, Wefel JS, Warren JJ. Acidic beverages increase the risk of in vitro tooth erosion. *Nutr Res.* (2008), 28(5):299-303.
- Ekshyyan VP, Hebert VY, Khandelwal A, Dugas TR. Resveratrol inhibits rat aortic vascular smooth muscle cell proliferation via estrogen receptor dependent nitric oxide production. *J Cardiovasc Pharmacol.* (2007), 50:83-93.
- Erdman JW Jr, Ford NA, Lindshield BL. Are the health attributes of lycopene related to its antioxidant function? *Arch Biochem Biophys.* (2009) 483:229-235
- Faith MS, Dennison BA, Edmunds LS, Stratton HH. Fruit juice intake predicts increased adiposity gain in children from low-income families: weight status-by environment interaction. *Pediatrics.* (2006), 118:2066-2075.
- Fardet A, Llorach R, Orsoni A, Martin JF, Pujos-Guillot E, Lapierre C, Scalbert A. Metabolomics provide new insight on the metabolism of dietary phytochemicals in rats. *J Nutr.* (2008), 138:1282-1287.
- Ferrazzano GF, Amato I, Ingenito A, De Natale A, Pollio A. Anti-cariogenic effects of polyphenols from plant stimulant beverages (cocoa, coffee, tea). *Fitoterapia.* (2009), 80(5):255-62.
- Foster-Powell K, Holt SA, Brand-Miller JC. International Table of Glycemic Index and Glycemic Load Values. *Am J Clin Nutr.* (2001), 76:5-56.
- Freedman JE, Parker C 3rd, Li L, Perlman JA, Frei B, Ivanov V, Deak LR, Iafrafi MD, Folts JD. Select flavonoids and whole juice from purple grapes inhibit platelet function and enhance nitric oxide release. *Circulation.* (2001),103:2792-2798.
- Frikke-Schmidt H, Lykkesfeldt J. Role of marginal vitamin C deficiency in atherogenesis: in vivo models and clinical studies. *Basic Clin Pharmacol Toxicol.* (2009),104:419-433.
- Fruit and vegetables for health: Report of a Joint FAO/WHO Workshop, (2004), Kobe, Japan.
- George TW, Niwat C, Waroonphan S, Gordon MH, Lovegrove JA. Effects of chronic and acute consumption of fruit- and vegetable-puree-based drinks on vasodilation, risk factors for CVD and the response as a result of the eNOS G298T polymorphism. *Proc Nutr Soc.* (2009), 68:148-61.

- Gerhaeuser C. Cancer chemopreventive potential of apples, apple juice, and apple components. *Planta Med.* (2008), 74:1608-1624.
- Ghanim H, Mohanty P, Pathak R, Chaudhuri A, Sia CL, Dandona P. Orange juice or fructose intake does not induce oxidative and inflammatory response. *Diabetes Care.* (2007), 30:1406-1411.
- Gonda TA, Tu S, Wang TC. Chronic inflammation, the tumor microenvironment and carcinogenesis. *Cell Cycle.* (2009), 8:2005-2013.
- González-Gallego J, García-Mediavilla MV, Sánchez-Campos S, Tuñón MJ. Fruit polyphenols, immunity and inflammation. *Br J Nutr.* (2010)104 Suppl 3:S15-27
- Gorinstein S, Caspi A, Libman I, Katrich E, Lerner HT, Trakhtenberg S. Fresh israeli jaffa sweetie juice consumption improves lipid metabolism and increases antioxidant capacity in hypercholesterolemic patients suffering from coronary artery disease: studies in vitro and in humans and positive changes in albumin and fibrinogen fractions. *J Agric Food Chem.* (2004), 52:5215-5222.
- Gorinstein S, Caspi A, Libman I, Lerner HT, Huang D, Leontowicz H, Leontowicz M, Tashma Z, Katrich E, Feng S, Trakhtenberg S. Red grapefruit positively influences serum triglyceride level in patients suffering from coronary atherosclerosis: studies in vitro and in humans. *J Agric Food Chem.* (2006), 54:1887-1892.
- Granic I, Dolga AM, Nijholt IM, van Dijk G, Eisel UL. Inflammation and NF-kappaB in Alzheimer's disease and diabetes. *J Alzheimers Dis.* (2009), 16:809-821.
- Grassi D, Desideri G, Croce G, Tiberti S, Aggio A, Ferri C. Flavonoids, vascular function and cardiovascular protection. *Curr Pharm Des.* (2009), 15:1072-1084.
- Guay DR. Cranberry and urinary tract infections. *Drugs.* (2009), 69:775-807.
- Habauzit V, Trzeciakiewicz A, Gil-izquierdo A, Mardon J, Morand C, Davicco MJ, Lebeque P, Offord E, Coxam V, Horcajada MN. Hesperidin and naringin, two main flavonoids of citrus fruits, improve bone metabolism in senescent male rats. *Third International Conference on Polyphenols and Health* (2007).
- Habauzit V, Nielsen IL, Gil-Izquierdo A, Trzeciakiewicz A, Morand C, Chee W, Barron D, Lebecque P, Davicco MJ, Williamson G, Offord E, Coxam V, Horcajada MN. Increased bioavailability of hesperetin-7-glucoside compared with hesperidin results in more efficient prevention of bone loss in adult ovariectomised rats. *Br J Nutr.* (2009), 27:1-9.
- Hadipour-Jahromy M, Mozaffari-Kermani R. Chondroprotective effects of pomegranate juice on monoiodoacetate-induced osteoarthritis of the knee joint of mice. *Phytother Res.* (2010), 24 (2): 182-185.
- Hannig C, Sorg J, Spitzmüller B, Hannig M, Al-Ahmad A. Polyphenolic beverages reduce initial bacterial adherence to enamel in situ. *J Dent.* (2009), 37(7):560-566.
- Hardcastle AC, Aucott L, Reid DM, Macdonald HM. Associations between dietary flavonoid intakes and bone health in a Scottish population. *J Bone Miner Res.* (2010) DOI: 10.1002/jbmr.285

- Hardy R, Cooper MS. Bone loss in inflammatory disorders. *J Endocrinol.* (2009), 201:309-320.
- Harrison FE, May JM. Vitamin C function in the brain: vital role of the ascorbate transporter SVCT2. *Free Radic Biol Med.* (2009), 46:719-730.
- Hochman N, Houry-Haddad Y, Koblinski J, Wahl L, Roniger M, Bar-Sinai A, Weiss EI, Hochman J. Cranberry juice constituents impair lymphoma growth and augment the generation of antilymphoma antibodies in syngeneic mice. *Nutr Cancer.* (2008), 60):511-517.
- Holst B, Williamson G. Nutrients and phytochemicals: from bioavailability to bioefficacy beyond antioxidants. *Curr Opin Biotechnol.* (2008),19(2):73-82.
- Holt EM, Steffen LM, Moran A, Basu S, Steinberger J, Ross JA, Hong CP, Sinaiko AR. Fruit and vegetable consumption and its relation to markers of inflammation and oxidative stress in adolescents. *J Am Diet Assoc.* (2009), 109:414-421.
- Hooper SM, Newcombe RG, Faller R, Eversole S, Addy M, West NX. The protective effects of toothpaste against erosion by orange juice: studies in situ and in vitro. *J Dent.* (2007), 35:476-481.
- Horcajada MN, Habauzit V, Trzeciakiewicz A, Morand C, Gil-Izquierdo A, Mardon J, Lebecque P, Davicco MJ, Chee WS, Coxam V, Offord E. Hesperidin inhibits ovariectomized-induced osteopenia and shows differential effects on bone mass and strength in young and adult intact rats. *J Appl Physiol.* (2008), 104:648-54.
- [http://www.efsa.europa.eu/EFSA/efsa\\_locale-1178620753812\\_article13.htm](http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_article13.htm) (2009)(database with list of health claims, art13claims.mdb)
- <http://www.fda.gov/Food/GuidanceComplianceRegulatoryInformation/GuidanceDocuments/FoodLabelingNutrition/FoodLabelingGuide/ucm064919.htm> (2008)
- <http://www.fruitsandveggiesmatter.gov>
- <http://www.mypyramid.gov/pyramid/fruits.html>.
- <http://www.phenol-explorer.eu/> (2009)
- Iriti M, Faoro F. Bioactivity of grape chemicals for human health. *Nat Prod Commun.* (2009), 4:611-634.
- Joseph JA, Shukitt-Hale B, Willis LM. Grape Juice, Berries, and Walnuts Affect Brain Aging and Behavior. *J Nutr.* (2009), 139:1813S-1817S.
- Kanapuru B, Ershler WB. Inflammation, coagulation, and the pathway to frailty. *Am J Med.* (2009)122:605-13
- Kannappan S, Anuradha CV. Naringenin enhances insulin-stimulated tyrosine phosphorylation and improves the cellular actions of insulin in a dietary model of metabolic syndrome. *Eur J Nutr.* 2009 Sep 2. [Epub ahead of print]
- Kelly RP, Poo Yeo K, Isaac HB, Lee CY, Huang SH, Teng L, Halliwell B, Wise SD. Lack of effect of acute oral ingestion of vitamin C on oxidative stress, arterial stiffness or blood pressure in healthy subjects. *Free Radic Res.* (2008) 42:514-522.
- King GL. The role of inflammatory cytokines in diabetes and its complications. *J Periodontol.* (2008),79 :Suppl1527-1534.

- Koltover V. Bioantioxidants: the systems reliability standpoint. *Toxicol Ind Health*. (2009), 25:295-299.
- Kyle JA, Sharp L, Little J, Duthie GG, McNeill G. Dietary flavonoid intake and colorectal cancer: a case-control study. *Br J Nutr*. 2009 7:1-8.
- Lam CK, Zhang Z, Yu H, Tsang SY, Huang Y, Chen ZY. Apple polyphenols inhibit plasma CETP activity and reduce the ratio of non-HDL to HDL cholesterol. *Mol Nutr Food Res*. (2008), 52:950–958.
- Landon S. Fruit juice nutrition and health. *Food Australia*. (2007), 59:533-538.
- Lee CY, Isaac HB, Huang SH, Long LH, Wang H, Gruber J, Ong CN, Kelly RP, Halliwell B. Limited antioxidant effect after consumption of a single dose of tomato sauce by young males, despite a rise in plasma lycopene. *Free Radic Res*. (2009), 43:622-628.
- Leifert WR, Abeywardena MY. Cardioprotective actions of grape polyphenols. *Nutr Res*. (2008), 28:729-737.
- Levine M, Dhariwal R K, Washko PW, Welch RW, Wang Y. Cellular functions of ascorbic acid: a means to determine vitamin C requirements. *Asia Pacific J Clin Nutr*. (1993), 2, Suppl 1, 5-13.
- Lim S, Sohn W, Burt BA, Sandretto AM, Kolker JL, Marshall TA, Ismail AI. Cariogenicity of soft drinks, milk and fruit juice in low-income african-american children: a longitudinal study. *J Am Dent Assoc*. (2008), 139:959-67.
- Macready AL, Kennedy OB, Ellis JA, Williams CM, Spencer JP, Butler LT. Flavonoids and cognitive function: a review of human randomized controlled trial studies and recommendations for future studies. *Genes Nutr*. (2009) 4(4):227-42.
- Maeng HG, Lim H, Jeong YJ, Woo A, Kang JS, Lee WJ, Hwang YI. Vitamin C enters mouse T cells as dehydroascorbic acid in vitro and does not recapitulate in vivo vitamin C effects. *Immunobiology*. (2009), 214:311-320.
- Mandadi K, Ramirez M, Jayaprakasha GK, Faraji B, Lihono M, Deyhim F, Patil BS. Citrus bioactive compounds improve bone quality and plasma antioxidant activity in orchidectomized rats. *Phytomedicine*. (2009),16:513-520.
- Manners GD. limonoids: analysis, bioactivity, and biomedical prospects. *J Agric Food Chem*. (2007), 55:8285-8294.
- Marques FZ, Markus MA, Morris BJ. Resveratrol: Cellular actions of a potent natural chemical that confers a diversity of health benefits. *Int J Biochem Cell Biol*. (2009), 41 :2125-2128.
- Martí N, Mena P, Cánovas JA, Micol V, Saura D. Vitamin C and the role of citrus juices as functional food. *Nat Prod Commun*. 2009, 4:677-700.
- Matsunaga N, Tsuruma K, Shimazawa M, Yokota S, Hara H. Inhibitory actions of bilberry anthocyanidins on angiogenesis. *Phytother Res*. (2010); 24 Suppl 1:S42-7
- Mattiello T, Trifirò E, Jotti GS, Pulcinelli FM. Effects of pomegranate juice and extract polyphenols on platelet function. *J Med Food*. (2009), 12:334-339.
- Mauray A, Felgines C, Morand C, Mazur A, Scalbert A, Milenkovic D. Nutrigenomic analysis of the protective effects of bilberry anthocyanin-rich extract in apo E-deficient mice *Genes Nutr*. (2010);5:343-353

- Mendosa R. New International Table of Glycemic Index (GI) and Glycemic Load Values. *Am J Clin Nutr.* (2003 ) 77(4):994
- Morand C, Milenkovic D., Lioger D., Martin JF, Dubray C, Scalbert A., Mazur A. Hesperidin contributes to the vascular protective effects of orange juice: a randomized crossover study in healthy. *Am J Clin Nutr.* (2011a), 93: 73-80
- Morand C., Milenkovic D., Chanet A., Deval C, Mazur A (2011b) Nutrigenomic effects of hesperidin, the major polyphenol of orange, related to its cardiovascular protective effects Experimental Biology 2011 Conference Washington, D.C
- Morin B, Nichols LA, Zalasky KM, Davis JW, Manthey JA, Holland LJ. The citrus flavonoids hesperetin and nobiletin differentially regulate low density lipoprotein receptor gene transcription in HepG2 liver cells. *J Nutr.* (2008), 138:1274-1281.
- Mori-Okamoto J, Otawara-Hamamoto Y, Yamato H, Yoshimura H. Pomegranate extract improves a depressive state and bone properties in menopausal syndrome model ovariectomized mice. *J Ethnopharmacol.* (2004), 92:93-101.
- Mulvihill EE, Allister EM, Sutherland BG, Telford DE, Sawyez CG, Edwards JY, Markle JM, Hegele RA, Huff MW. Naringenin prevents dyslipidemia, apoB overproduction and hyperinsulinemia in LDL-receptor null mice with diet-induced insulin resistance. *Diabetes.* (2009) 58(10):2198-210.
- Murakami A, Song M, Katsumata S, Uehara M, Suzuki K, Ohigashi H. Citrus nobiletin suppresses bone loss in ovariectomized ddY mice and collagen-induced arthritis in DBA/1J mice: possible involvement of receptor activator of NF-kappaB ligand (RANKL)-induced osteoclastogenesis regulation. *Biofactors.* (2007), 30:179-192.
- Nahmias Y, Goldwasser J, Casali M, van Poll D, Wakita T, Chung RT, Yarmush ML. Apolipoprotein B-dependent hepatitis C virus secretion is inhibited by the grapefruit flavonoid naringenin. *Hepatology.* (2008), 47:1437-1445.
- Neto CC. Cranberry and blueberry: evidence for protective effects against cancer and vascular diseases. *Mol Nutr Food Res.* (2007) ,51:652-664.
- Neukam K, Stahl W, Tronnier H, Sies H, Heinrich U. Consumption of flavanol-rich cocoa acutely increases microcirculation in human skin. *Eur J Nutr.* (2007), 46:53-56.
- Newby PK, Peterson KE, Berkey CS, Leppert J, Willett WC, Colditz GA. Beverage consumption is not associated with changes in weight and body mass index among low-income preschool children in North Dakota. *J Am Diet Assoc.* (2004), 104:1086-1094.
- Nicklas TA, O'Neil CE, Kleinman R. Association between 100% juice consumption and nutrient intake and weight of children aged 2 to 11 years. *Arch Pediatr Adolesc Med.* (2008), 162:557-565.
- O'Neil, CE and Nicklas, TA. A Review of the Relationship Between 100% Fruit Juice Consumption and Weight in Children and Adolescents. *American Journal of Lifestyle Medicine.* (2008), 2:315-354.
- O'Connor TM, Yang SJ, Nicklas TA. Beverage intake among preschool children and its effect on weight status. *Pediatrics.* (2006), 118:e1010-1018.

- Ovesná J, Slabý O, Toussaint O, Kodíček M, Marsík P, Pouchová V, Vaněk T. High throughput 'omics' approaches to assess the effects of phytochemicals in human health studies. *Br J Nutr.* (2008),99E Suppl: ES127-134.
- Palmer JR, Boggs DA, Krishnan S, Hu FB, Singer M, Rosenberg L. Sugar-Sweetened Beverages and Incidence of Type 2 Diabetes Mellitus in African American Women. *Arch Int Med.* (2008), 168:1487-1492.
- Pezzuto JM, Venkatasubramanian V, Hamad M, Morris KR. Unraveling the Relationship between Grapes and Health. *J Nutr.* (2009),139:1783S-1787S.
- Poulose SM, Harris ED, Patil BS. Antiproliferative effects of citrus limonoids against human neuroblastoma and colonic adenocarcinoma cells. *Nutr Cancer.* (2006), 56:103-112.
- Qin Y, Xia M, Ma J, Hao Y, Liu J, Mou H, Cao L, Ling W. Anthocyanin supplementation improves serum LDL- and HDL-cholesterol concentrations associated with the inhibition of cholesteryl ester transfer protein in dyslipidemic subjects. *Am J Clin Nutr.* (2009), 90:485-492
- Ren YF, Amin A, Malmstrom H. Effects of tooth whitening and orange juice on surface properties of dental enamel. *J Dent.* (2009), 37:424-31.
- Reshef N, Hayari Y, Goren C, Boaz M, Madar Z, Knobler H. Antihypertensive effect of sweetie fruit in patients with stage I hypertension. *Am J Hypertens.* (2005), 18:1360-1363.
- Ross SM. Pomegranate: its role in cardiovascular health. *Holist Nurs Pract.* (2009), 23:195-197.
- Ruxton CH, Gardner EJ, Walker D. Can pure fruit and vegetable juices protect against cancer and cardiovascular disease too? A review of the evidence. *Int J Food Sci Nutr.* (2006), 57:249-272.
- Sahni S, Hannan MT, Blumberg J, Cupples LA, Kiel DP, Tucker KL. Protective effect of total carotenoid and lycopene intake on the risk of hip fracture: a 17-year follow-up from the Framingham Osteoporosis Study. *J Bone Miner Res.* (2009a), 24:1086-1094.
- Sahni S, Hannan MT, Blumberg J, Cupples LA, Kiel DP, Tucker KL. Inverse association of carotenoid intakes with 4-y change in bone mineral density in elderly men and women: the Framingham Osteoporosis Study. *Am J Clin Nutr.* (2009b) 89:416-424.
- Sánchez-Moreno C, Jiménez-Escrig A, Martín A. Stroke: roles of B vitamins, homocysteine and antioxidants. *Nutr Res Rev.* (2009), 22:49-67.
- Scalbert A, Johnson IT, Saltmarsh M. Polyphenols: antioxidants and beyond. *Am J Clin Nutr.* (2005), 81:215S-217S.
- Scalbert A, Knasmüller S. Genomic effects of phytochemicals and their implication in the maintenance of health. *J Nutr.* (2008), 99E Suppl ES1-2.
- Seeram NP. Berry fruits: compositional elements, biochemical activities, and the impact of their intake on human health, performance, and disease. *J Agric Food Chem.* (2008), 56:627-629.
- Serafini M. Rise and fall of dietary antioxidants for disease prevention: magic bullets; false myth or scare knowledge. *Fav health.* (2009).

- Sheretz E, Goldsmith LA. Nutritional influences on the skin. In: Goldsmith LA, ed. *Physiology, biochemistry, and molecular biology of the skin*. New York, NY Oxford University Press. (1991), 1315–1328.
- Shukitt-Hale B, Cheng V, Joseph JA. Effects of blackberries on motor and cognitive function in aged rats. *Nutr Neurosci*. (2009), 12:135-140.
- Shukitt-Hale B, Lau FC, Joseph JA. Berry fruit supplementation and the aging brain. *J Agric Food Chem*. (2008),56:636-641.
- Shukla M, Gupta K, Rasheed Z, Khan KA, Haqqi TM. Consumption of hydrolyzable tannins-rich pomegranate extract suppresses inflammation and joint damage in rheumatoid arthritis. *Nutrition*. (2008), 24:733-743.
- Sies H. Total antioxidant capacity: appraisal of a concept. *J Nutr*. (2007), 137:1493-1495.
- Skinner JD, Carruth BR, Moran J, Houck K, and Coletta F. Fruit Juice Intake Is Not Related to Children's Growth. *Pediatrics*. (1999), 103:58-64.
- Skinner JD, Carruth BR. A longitudinal study of children's juice intake and growth: the juice controversy revisited. *J Am Diet Assoc*. (2001), 101: 432–437.
- Soler C, Soriano JM, Mañes J. Apple-products phytochemicals and processing: a review. *Nat Prod Commun*. (2009), 4:659-670.
- Spencer JP. Flavonoids and brain health: multiple effects underpinned by common mechanisms. *Genes Nutr*. (2009b), *Genes Nutr*. (2009), 4(4):243-50
- Spencer JP. Flavonoids: modulators of brain function? *Br J Nutr*. (2008) 99 E Suppl 1:ES60-77
- Spencer JP. The impact of flavonoids on memory: physiological and molecular considerations. *Chem Soc Rev*. (2009a), 38:1152-61.
- Spencer JP. The impact of fruit flavonoids on memory and cognition *Br J. Nutr* (2010), 104, S40–S47
- Stahl W, Sies H. Carotenoids and flavonoids contribute to nutritional protection against skin damage from sunlight. *Mol Biotechnol*. (2007), 37:26-30.
- Steiner C, Arnould S, Scalbert A, Manach C. Isoflavones and the prevention of breast and prostate cancer: new perspectives opened by nutrigenomics. *Br J Nutr*. (2008), 99 E Suppl ES78-108.
- Stevenson DE, Hurst RD. Polyphenolic phytochemicals—just antioxidants or much more? *Cell Mol Life Sci*. (2007), 64:2900-2916.
- Stipevic T, Piljac J, Vanden Berghe D. Effect of different flavonoids on collagen synthesis in human fibroblasts. *Plant Foods Hum Nutr*. (2006), 61:29-34.
- Sugiura M, Nakamura M, Ogawa K, Ikoma Y, Ando F, Yano M. Bone mineral density in post-menopausal female subjects is associated with serum antioxidant carotenoids. *Osteoporos Int*. (2008), 19: 211-219.
- Szajdek A, Borowska EJ. Bioactive compounds and health-promoting properties of berry fruits: a review. *Plant Foods Hum Nutr*. (2008),63:147-156.
- Tanaka K, Miyake Y, Sasaki S, Ohya Y, Matsunaga I, Yoshida T, Hirota Y, Hajime O, Kanzaki H, Kitada M, Horikoshi Y, Ishiko O, Nakai Y, Nishio J, Yamamasu S,

- Yasuda J, Kawai S, Yanagihara K, Wakuda K, Kawashima T, Narimoto K, Iwasa Y, Orino K, Tsunetoh I, Yoshida J, Ito J, Kaneko T, Sanfujinka K, Kamiya T, Kuribayashi H, Taniguchi T, Takemura H, Morimoto Y. Beverage consumption and the prevalence of tooth loss in pregnant Japanese women: the Osaka Maternal and Child Health Study. Osaka Maternal and Child Health Study Group. *Fukuoka Igaku Zasshi*. (2008), 99:80-89.
- Thomas AE. Some observations on the influence of orange juice ingestion on the teeth and supporting structures. *Oral Surg Oral Med Oral Pathol*. (1954), 7:741-749.
- Thomasset S, Berry DP, Cai H, West K, Marczylo TH, Marsden D, Brown K, Dennison A, Garcea G, Miller A, Hemingway D, Steward WP, Gescher AJ. Pilot study of oral anthocyanins for colorectal cancer chemoprevention. *Cancer Prev Res (Phila Pa)*. (2009), 2:625-633.
- Trzeciakiewicz A, Habauzit V, Horcajada MN. When nutrition interacts with osteoblast function: molecular mechanisms of polyphenols. *Nutr Res Rev*. (2009), 22:68-81.
- Trzeciakiewicz A, Habauzit V, Mercier S, Lebecque P, Davicco MJ, Coxam V, Demigne C, Horcajada MN. Hesperetin stimulates differentiation of primary rat osteoblasts involving the BMP signalling pathway. *J Nutr Biochem*. (2010) 21(5):424-31
- Vafeiadou K, Vauzour D, Lee HY, Rodriguez-Mateos A, Williams RJ, Spencer JP. The citrus flavanone naringenin inhibits inflammatory signalling in glial cells and protects against neuroinflammatory injury. *Arch Biochem Biophys*. (2009), 484:100-109.
- Veeriah S, Miene C, Habermann N, Hofmann T, Klenow S, Sauer J, Böhmer F, Wölfl S, Pool-Zobel BL. Apple polyphenols modulate expression of selected genes related to toxicological defence and stress response in human colon adenoma cells. *Int J Cancer*. (2008)122:2647-2655.
- Venn BJ, Green TJ Glycemic index and glycemic load: measurement issues and their effect on diet-disease relationships. *Eur J Clin Nutr*. (2007) 61 Suppl 1:S122-131.
- Vrolix R, Mensink RP. Variability of the glycemic response to single food products in healthy subjects. *Contemp Clin Trials*. (2010 ) 31(1): 5-11.
- Wang J, Ho L, Zhao W, Ono K, Rosensweig C, Chen L, Humala N, Teplow DB, Pasinetti GM. Grape-derived polyphenolics prevent Abeta oligomerization and attenuate cognitive deterioration in a mouse model of Alzheimer's disease. *J Neurosci*. (2008),28:6388-8892.
- Wärnberg J, Gomez-Martinez S, Romeo J, Díaz LE, Marcos A. Nutrition, inflammation, and cognitive function. *Ann N Y Acad Sci*. (2009), 1153:164-175.
- Willershausen B, Callaway A, Azrak B, Duschner H. Influence of apple juice on human enamel surfaces of the first and second dentition - an in vitro study. *Eur J Med Res*. (2008), 13:349-354.
- Williams RJ, Spencer JP, Rice-Evans C. Flavonoids: antioxidants or signaling molecules? *Free Radic Biol Med*. (2004), 36:838-49.
- Williamson G, Holst B. Dietary reference intake (DRI) value for dietary polyphenols:are we heading in the right direction? *Br J Nutr*. (2008),99 Suppl

3:S55-58.

Willis LM, Shukitt-Hale B, Joseph JA. Recent advances in berry supplementation and age-related cognitive decline. *Curr Opin Clin Nutr Metab Care.* (2009), 12:91-94.

Wilson T, Meyers SL, Singh AP, Limburg PJ, Vorsa N. Favorable glycemic response of type 2 diabetics to low-calorie cranberry juice. *J Food Sci.* (2008),73:H241-245.

Wong RW, Rabie AB. Effect of naringin on bone cells. *J Orthop Res.* (2006),24:2045-2050.

Wu CH, Wu CF, Huang HW, Jao YC, Yen GC. Naturally occurring flavonoids attenuate high glucose-induced expression of proinflammatory cytokines in human monocytic THP-1 cells. *Mol Nutr Food Res.* (2009), 53:984-95.

Wu F, Schuster DP, Tysl K, Wilson JX. Ascorbate inhibits NADPH oxidase subunit p47phox expression in microvascular endothelial cells. *Free Radic Biol Med.* (2007), 42:124-131.

Wu H, Dai Q, Shrubsole MJ, Ness RM, Schlundt D, Smalley WE, Chen H, Li M, Shyr Y, Zheng W. Fruit and vegetable intakes are associated with lower risk of colorectal adenomas. *J Nutr.* (2009), 139:340-344.

Yao F, Shu G, Yang Y, Shi B. Grapefruit and diet of the type 2 Diabetes: 53 (2): A591 (2004).

Yoshida M, McKeown NM, Rogers G, Meigs JB, Saltzman E, D'Agostino R, Jacques PF. Surrogate markers of insulin resistance are associated with consumption of sugar-sweetened drinks and fruit juice in middle and older-aged adults. *J Nutr.* (2007), 137(9):2121-2127.

Youdim KA, Dobbie MS, Kuhnle G, Proteggente AR, Abbott NJ, Rice-Evans C. Interaction between flavonoids and the blood-brain barrier: in vitro studies. *J Neurochem.* (2003), 85:180-192.